

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date
29 January 2004 (29.01.2004)

PCT

(10) International Publication Number
WO 2004/010118 A1

(51) International Patent Classification⁷: G01N 21/31, 21/35, A61B 5/103

(74) Agent: DUNNETT, Julie, Elizabeth; Barker Brettell, St John's Innovation Centre, Cowley Road, Cambridge CB4 0WS (GB).

(21) International Application Number:
PCT/GB2003/003245

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 7 July 2003 (07.07.2003)

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language: English

Published:

(26) Publication Language: English

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

[Continued on next page]

(54) Title: METHOD AND APPARATUS FOR INVESTIGATING HISTOLOGY OF EPITHELIAL TISSUE

$$Z = \frac{R_j(\lambda_1)}{R_i(\lambda_1)^j} \quad (I)$$

$$Z = \frac{R_j(c, h, \lambda_1)^j}{R_i(c, h, \lambda_2)^j} = \frac{R_j(\lambda_1)^j}{R_i(\lambda_2)^j} = \frac{R_j(\lambda_1)}{R_i(\lambda_2)^j} \quad (II)$$



Invasive BCC with the Z image on the right showing marked dermal involvement

WO 2004/010118 A1

(57) Abstract: A method for monitoring the presence of selected chromophores in a sample of epithelial tissue, independent of the amount of a predetermined chromophore, the method comprising: illuminating an area of tissue by projecting light from a light source of at least two different wavelengths λ_1, λ_2 ; receiving light remitted by the illuminated area of tissue at a photoreceptor; analysing the received light to identify and measure the proportion of light of each wavelength remitted from the tissue $I_r(\lambda)$; calculating the ratio of light at each wavelength returned from the tissue $R_i(\lambda)$, and then calculating $Z = \text{Formula (I)}$; where i is chosen such that Z is independent of the amount of predetermined chromophore. Typically i is calculated such that $Z = \text{Formula (II)}$; where j and k are such that $2j\alpha(\lambda_1) = 2kj\alpha(\lambda_2) = 1$ where $\alpha(\lambda_1)$ and $\alpha(\lambda_2)$ are the absorption coefficients for the predetermined chromophore at each wavelength.

BEST AVAILABLE COPY